

This information has not been peer-reviewed. Responsibility for the findings rests solely with the author(s).

Deposited research article

## The functional genomic response of developing embryonic submandibular glands to NF $\kappa$ B inhibition

Michael Melnick, Haiming Chen, Yan Min Zhou and Tina Jaskoll

Address: Laboratory for Developmental Genetics, University of Southern California, Los Angeles, CA 90089-0641, USA.

Correspondence: Michael Melnick. E-mail: mmelnick@hsc.usc.edu

Posted: 4 May 2001

*Genome Biology* 2001, **2(6)**:preprint0005.1-0005.44

The electronic version of this article is the complete one and can be found online at <http://genomebiology.com/2001/2/6/preprint/0005>

© BioMed Central Ltd (Print ISSN 1465-6906; Online ISSN 1465-6914)

Received: 27 April 2001

This was the first version of this article to be made available publicly. The full version of this preprint has been removed at the request of the authors. A peer-reviewed and modified version of this manuscript is now available at <http://biomedcentral.com/1471-213X/1/15/>



deposited research

AS A SERVICE TO THE RESEARCH COMMUNITY, GENOME **BIOLOGY** PROVIDES A 'PREPRINT' DEPOSITORY TO WHICH ANY PRIMARY RESEARCH CAN BE SUBMITTED AND WHICH ALL INDIVIDUALS CAN ACCESS FREE OF CHARGE. ANY ARTICLE CAN BE SUBMITTED BY AUTHORS, WHO HAVE SOLE RESPONSIBILITY FOR THE ARTICLE'S CONTENT. THE ONLY SCREENING IS TO ENSURE RELEVANCE OF THE PREPRINT TO GENOME **BIOLOGY**'S SCOPE AND TO AVOID ABUSIVE, LIBELLOUS OR INDECENT ARTICLES. ARTICLES IN THIS SECTION OF THE JOURNAL HAVE **NOT** BEEN PEER REVIEWED. EACH PREPRINT HAS A PERMANENT URL, BY WHICH IT CAN BE CITED. RESEARCH SUBMITTED TO THE PREPRINT DEPOSITORY MAY BE SIMULTANEOUSLY OR SUBSEQUENTLY SUBMITTED TO GENOME **BIOLOGY** OR ANY OTHER PUBLICATION FOR PEER REVIEW; THE ONLY REQUIREMENT IS AN EXPLICIT CITATION OF, AND LINK TO, THE PREPRINT IN ANY VERSION OF THE ARTICLE THAT IS EVENTUALLY PUBLISHED. IF POSSIBLE, GENOME **BIOLOGY** WILL PROVIDE A RECIPROCAL LINK FROM THE PREPRINT TO THE PUBLISHED ARTICLE.



**The Functional Genomic Response of Developing  
Embryonic Submandibular Glands to NF $\kappa$ B Inhibition**

Michael Melnick, Haiming Chen, Yan Min Zhou and Tina Jaskoll

Laboratory for Developmental Genetics, University of Southern California

Address Correspondence to:

Dr. Michael Melnick

Laboratory for Developmental Genetic

University of Southern California

925 W 34<sup>th</sup> Street DEN 4266

Los Angeles, CA 90089-0641

[mmelnick@hsc.usc.edu](mailto:mmelnick@hsc.usc.edu)

213-740-1417 (Voice)

213-740-7560 (Fax)

## **Abstract**

**Background:** The proper balance between epithelial cell proliferation, quiescence, and apoptosis during development is mediated by the specific temporal and spatial appearance of transcription factors, growth factors, cytokines, caspases, etc. Since prior studies suggest the importance of transcription factor NF $\kappa$ B during embryonic submandibular salivary gland (SMG) development, we attempted to delineate the emergent dynamics of a cognate signaling network by studying the molecular patterns and phenotypic outcomes of interrupted NF $\kappa$ B signaling in embryonic SMG explants.

**Results:** SN50-mediated inhibition of NF $\kappa$ B signaling in SMG explants results in a highly significant increase in apoptosis and decrease in cell proliferation. Probabilistic Neural Network (PNN) analyses of proteomic and transcriptomic assays identify specific transcripts and proteins with altered expression that best discriminate control from SN50-treated SMGs with 100% sensitivity and specificity. These include PCNA, GR, BMP1, BMP3b, Chk1, Caspase 6, E2F1, c-Raf, and JNK-1, as well as several others of lesser importance. Thus, it is apparent that inhibition of NF $\kappa$ B nuclear translocation alters the expression of a cognate genetic network with broadly related, rather than independent, components.

**Conclusions:** Morphological and functional genomic analyses subsequent to inhibition of NF $\kappa$ B nuclear translocation indicate that NF $\kappa$ B-mediated transcription is critical to embryonic SMG developmental homeostasis. Relative to understanding complex genetic networks and organogenesis, our results demonstrate the importance of evaluating the gene, protein, and activated protein expression of multiple components from multiple pathways within broad functional categories.